

A two armed bandit type problem revisited

GILLES PAGÈS *

February 1, 2008

Abstract

In [2] M. Benaïm and G. Ben Arous solve a multi-armed bandit problem arising in the theory of learning in games. We propose an short elementary proof of this result based on a variant of the Kronecker Lemma.

Key words: Two-armed bandit problem, Kronecker Lemma, learning theory, stochastic fictitious play.

In [2] a multi-armed bandit problem is addressed and investigated by M. Benaïm and G. Ben Arous. Let $f_0, \dots, f_d$ denote $d + 1$ real-valued continuous functions defined on $[0, 1]^{d+1}$. Given a sequence $x = (x_n)_{n \geq 1} \in \{0, \dots, d\}^{\mathbb{N}^*}$ (the *strategy*), set for every $n \geq 1$

$$\bar{x}_n := (\bar{x}_n^0, \bar{x}_n^1, \dots, \bar{x}_n^d) \quad \text{with} \quad \bar{x}_n^i := \frac{1}{n} \sum_{k=1}^n \mathbf{1}_{\{x_k=i\}}, \quad i = 0, \dots, d,$$

and

$$Q(x) = \liminf_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=0}^{n-1} f_{x_{k+1}}(\bar{x}_k).$$

($\bar{x}_0 := (\bar{x}_0^0, \bar{x}_0^1, \dots, \bar{x}_0^d) \in [0, 1]^{d+1}$, $\bar{x}_0^0 + \dots + \bar{x}_0^d = 1$ is a starting distribution). Imagine $d + 1$ players enrolled in a cooperative/competitive game with the following simple rules: if player $i \in \{0, \dots, d\}$ plays at time n he is rewarded by $f_i(\bar{x}_n)$, otherwise he gets nothing; only one player can play at the same time. Then the sequence x is a playing strategy for the group of players and $Q(x)$ is the *global* cumulative worst payoff rate of the strategy x for the whole community of players (regardless of the cumulative payoff rate of each player).

In [2] an answer (see Theorem 1 below) is provided to the following question

What are the good strategies (for the group) ?

*Laboratoire de Probabilités et Modélisation aléatoire, UMR 7599, Université Paris 6, case 188, 4, pl. Jussieu, F-75252 Paris Cedex 5. E-mail: gpa@ccr.jussieu.fr

The authors rely on some recent tools developed in stochastic approximation theory (see e.g. [1]). The aim of this note is to provide an elementary and shorter proof based on a slight improvement of the Kronecker Lemma.

Let $\mathcal{S}_d := \{v \in [0, 1]^d, \sum_{i=1}^d v_i \leq 1\}$ and $\mathcal{P}_{d+1} := \{u \in [0, 1]^{d+1}, \sum_{i=1}^{d+1} u_i = 1\}$. Furthermore, for notational convenience, set

$$\begin{aligned} \forall v = (v_1, \dots, v_d) \in \mathcal{S}_d, \quad \tilde{v} &:= (1 - \sum_{i=1}^d v_i, v_1, \dots, v_d) \in \mathcal{P}_{d+1}, \\ \forall u = (u_0, u_1, \dots, u_d) \in \mathcal{P}_{d+1}, \quad \hat{u} &:= (u_1, \dots, u_d) \in \mathcal{S}_d. \end{aligned}$$

The canonical inner product on $\mathbb{R}^d$ will be denoted by $(v|v') = \sum_{i=1}^d v_i v'_i$. The interior of a subset A of $\mathbb{R}^d$ will be denoted $\overset{\circ}{A}$. For a sequence $u = (u_n)_{n \geq 1}$, $\Delta u_n := u_n - u_{n-1}$, $n \geq 1$.

The main result is the following theorem (first established in [2]).

Theorem 1 *Assume there is a function $\Phi : \mathcal{S}_d \rightarrow \mathbb{R}$, continuously differentiable on $\overset{\circ}{\mathcal{S}}_d$ having a continuous extension $\nabla \Phi$ on $\mathcal{S}_d$ and satisfying:*

$$\forall v \in \mathcal{S}_d, \quad \nabla \Phi(v) = (f_i(\tilde{v}) - f_0(\tilde{v}))_{1 \leq i \leq d}. \quad (1)$$

Set for every $u \in \mathcal{P}_{d+1}$,

$$q(u) := \sum_{i=0}^{d+1} u_i f_i(u)$$

and $Q^* := \max \{q(u), u \in \mathcal{P}_{d+1}\}$. Then, for every strategy $x \in \{0, 1, \dots, d\}^{\mathbb{N}^*}$,

$$Q(x) \leq Q^*.$$

Furthermore, for any strategy x such that $\bar{x}_n \rightarrow \bar{x}_\infty$,

$$\frac{1}{n} \sum_{k=1}^n f_{x_{k+1}}(\bar{x}_k) \rightarrow q(\bar{x}_\infty) \quad \text{as } n \rightarrow \infty \quad (\text{so that } Q(x) = q(\bar{x}_\infty)).$$

In particular there is no better strategy than choosing the player at random according to an i.i.d. strategy with distribution $\bar{x}^* \in \operatorname{argmax} q$.

The key of the proof is the following slight extension of the Kronecker Lemma.

Lemma 1 (“à la Kronecker” Lemma) *Let $(b_n)_{n \geq 1}$ be a nondecreasing sequence of positive real numbers converging to $+\infty$ and let $(a_n)_{n \geq 1}$ be a sequence of real numbers. Then*

$$\liminf_{n \rightarrow +\infty} \sum_{k=1}^n \frac{a_k}{b_k} \in \mathbb{R} \quad \implies \quad \liminf_{n \rightarrow +\infty} \frac{1}{b_n} \sum_{k=1}^n a_k \leq 0.$$

Proof. Set $C_n = \sum_{k=1}^n \frac{a_k}{b_k}$, $n \geq 1$ and $C_0 = 0$ so that $a_n = b_n \Delta C_n$. As a consequence, an Abel transform yields

$$\begin{aligned} \frac{1}{b_n} \sum_{k=1}^n a_k &= \frac{1}{b_n} \sum_{k=1}^n b_k \Delta C_k = \frac{1}{b_n} \left(b_n C_n - \sum_{k=1}^n C_{k-1} \Delta b_k \right) \\ &= C_n - \frac{1}{b_n} \sum_{k=1}^n C_{k-1} \Delta b_k. \end{aligned}$$

Now, $\liminf_{n \rightarrow +\infty} C_n$ being finite, for every $\varepsilon > 0$, there is an integer n_ε such that for every $k \geq n_\varepsilon$, $C_k \geq \liminf_{n \rightarrow +\infty} C_n - \varepsilon$. Hence

$$\frac{1}{b_n} \sum_{k=1}^n C_{k-1} \Delta b_k \geq \frac{1}{b_n} \sum_{k=1}^{n_\varepsilon} C_{k-1} \Delta b_k + \frac{b_n - b_{n_\varepsilon}}{b_n} \left(\liminf_k C_k - \varepsilon \right).$$

Consequently, $\liminf_{n \rightarrow +\infty} C_n$ being finite, one concludes that

$$\liminf_{n \rightarrow +\infty} \frac{1}{b_n} \sum_{k=1}^n a_k \leq \liminf_{n \rightarrow +\infty} C_n - 0 - 1 \times \left(\liminf_{k \rightarrow +\infty} C_k - \varepsilon \right) = \varepsilon. \quad \diamond$$

Proof of Theorem 1. First note that for every $u = (u_0, \dots, u_d) \in \mathcal{P}_{d+1}$,

$$q(u) := \sum_{i=0}^{d+1} u_i f_i(u) = f_0(u) + \sum_{i=1}^d u_i (f_i(u) - f_0(u))$$

so that

$$Q^* = \sup_{v \in \mathcal{S}_d} \left\{ f_0(\tilde{v}) + \sum_{i=1}^d v_i (f_i(\tilde{v}) - f_0(\tilde{v})) \right\} = \sup_{v \in \mathcal{S}_d} \{ f_0(\tilde{v}) + (v | \nabla \Phi(v)) \}.$$

Now, for every $k \geq 0$

$$\begin{aligned} f_{x_{k+1}}(\bar{x}_k) - q(\bar{x}_k) &= \sum_{i=0}^d (f_i(\bar{x}_k) \mathbf{1}_{\{x_{k+1}=i\}} - \bar{x}_k^i f_i(\bar{x}_k)) = \sum_{i=0}^d f_i(\bar{x}_k) (\mathbf{1}_{\{x_{k+1}=i\}} - \bar{x}_k^i) \\ &= \sum_{i=0}^d f_i(\bar{x}_k) (k+1) \Delta \bar{x}_{k+1}^i \\ &= (k+1) \sum_{i=1}^d (f_i(\bar{x}_k) - f_0(\bar{x}_k)) \Delta \bar{x}_{k+1}^i. \end{aligned}$$

The last equality reads using Assumption (1),

$$f_{x_{k+1}}(\bar{x}_k) - q(\bar{x}_k) = (k+1) (\nabla \Phi(\hat{\hat{x}}_k) | \Delta \hat{\hat{x}}_{k+1})$$

Consequently, by the fundamental formula of calculus applied to Φ on $(\hat{x}_k, \hat{x}_{k+1}) \subset \mathring{\mathcal{S}}_d$,

$$\begin{aligned} \frac{1}{n} \sum_{k=0}^{n-1} f_{x_{k+1}}(\bar{x}_k) - q(\bar{x}_k) &= \frac{1}{n} \sum_{k=0}^{n-1} (k+1) (\Phi(\hat{x}_{k+1}) - \Phi(\hat{x}_k)) - R_n \\ \text{with} \quad R_n &:= \frac{1}{n} \sum_{k=0}^{n-1} (\nabla \Phi(\hat{\xi}_k) - \nabla \Phi(\hat{x}_k) | (k+1) \Delta \hat{x}_{k+1}) \end{aligned}$$

and $\hat{\xi}_k \in (\hat{x}_k, \hat{x}_{k+1})$, $k = 1, \dots, n$. The fact that $|(k+1) \Delta \hat{x}_{k+1}| \leq 1$ implies

$$|R_n| \leq \frac{1}{n} \sum_{k=0}^{n-1} w(\nabla \Phi, |\Delta \hat{x}_{k+1}|)$$

where $w(g, \delta)$ denotes the uniform continuity δ -modulus of a function g . One derives from the uniform continuity of $\nabla \Phi$ on the compact set $\mathcal{S}_d$ that

$$R_n \rightarrow 0 \quad \text{as} \quad n \rightarrow +\infty.$$

Finally, the continuous function Φ being bounded on the compact set $\mathcal{S}_d$, the partial sums

$$\sum_{k=0}^{n-1} \Phi(\hat{x}_{k+1}) - \Phi(\hat{x}_k) = \Phi(\hat{x}_{n+1}) - \Phi(\hat{x}_0)$$

remain bounded as n goes to infinity. Lemma 1 then implies that

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=0}^{n-1} (k+1) (\Phi(\hat{x}_{k+1}) - \Phi(\hat{x}_k)) \leq 0.$$

One concludes by noting that on one hand

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} q(\bar{x}_k) \leq Q^* = \sup_{\mathcal{P}_{d+1}} q$$

and that, on the other hand, the function q being continuous,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} q(\bar{x}_k) = q(x^*) \quad \text{as soon as} \quad \bar{x}_n \rightarrow x^*. \quad \diamond$$

Corollary 1 *When $d+1 = 2$ (two players), Assumption (1) is satisfied as soon as f_0 and f_1 are continuous on $\mathcal{P}_2$ and then the conclusions of Theorem 1 hold true.*

Proof: This follows from the obvious fact that the continuous function $u_1 \mapsto f_1(1 - u_1, u_1) - f_0(1 - u_1, u_1)$ on $[0, 1]$ has an antiderivative. $\diamond$

FURTHER COMMENTS: • If one considers a slightly more general game in which some *weighted strategies* are allowed, the final result is not modified in any way provided the

weight sequence satisfies a very light assumption. Namely, assume that at time n the reward is

$$\Delta_{n+1} f_{x_{n+1}}(\bar{x}_n) \quad \text{instead of} \quad f_{x_{n+1}}(\bar{x}_n)$$

where the weight sequence $\Delta = (\Delta_n)_{n \geq 1}$ satisfies

$$\Delta_n \geq 0, \quad n \geq 1, \quad S_n = \sum_{k=1}^n \Delta_k \rightarrow +\infty, \quad \frac{\Delta_n}{S_n} \rightarrow 0 \text{ as } n \rightarrow \infty$$

then the quantities $\bar{x}_0^\Delta \in \mathcal{P}_{d+1}$, $\bar{x}_n^\Delta := (\bar{x}_n^{\Delta,0}, \dots, \bar{x}_n^{\Delta,d})$ with $\bar{x}_n^{\Delta,i} = \frac{1}{S_n} \sum_{k=1}^n \Delta_k \mathbf{1}_{\{x_k=i\}}$, $i = 0, \dots, d$, $n \geq 1$, and $Q^\Delta(x) = \liminf_{n \rightarrow +\infty} \frac{1}{S_n} \sum_{k=0}^{n-1} \Delta_{k+1} f_{x_{k+1}}(\bar{x}_k^\Delta)$ satisfy all the conclusions of Theorem 1 *mutatis mutandis*.

• Several applications of Theorem 1 to the theory of learning in games and to stochastic fictitious play are extensively investigated in [2] which we refer to for all these aspects. As far as we are concerned we will simply make a remark about some “natural” strategies which illustrates the theorem in an elementary way.

In the reward function at time k , *i.e.* $f_{x_k}(\bar{x}_{k-1})$, x_k represents the competitive term (“who will play ?”) and $\bar{x}_{k-1}$ represents a cooperative term (everybody’s past behaviour has influence on everybody’s reward).

This cooperative/competitive antagonism induces that in such a game a *greedy* competitive strategy is usually not optimal (when the players do not play a symmetric rôle). Let us be more specific. Assume for the sake of simplicity that $d+1 = 2$ (two players). Then one may consider without loss of generality that $\bar{x}_n = \hat{x}_n$ *i.e.* that $\bar{x}_n$ is a $[0, 1]$ -valued real number. A *greedy competitive* strategy is defined by

$$\text{player 1 plays at time } n \text{ (i.e. } x_n = 1) \text{ iff } f_1(\bar{x}_{n-1}) \geq f_0(\bar{x}_{n-1}) \quad (2)$$

i.e. the player with the highest reward is nominated to play. Note that such a strategy is anticipative from a probabilistic viewpoint. Then, for every $n \geq 1$,

$$f_{x_n}(\bar{x}_{n-1}) = \max(f_0(\bar{x}_{n-1}), f_1(\bar{x}_{n-1}))$$

and it is clear that

$$f_{x_n}(\bar{x}_{n-1}) - q(\bar{x}_n) = \max(f_0(\bar{x}_{n-1}), f_1(\bar{x}_{n-1})) - q(\bar{x}_n) =: \varphi(\bar{x}_n) \geq 0.$$

On the other hand, the proof of Theorem 1 implies that

$$\liminf_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=0}^{n-1} \varphi(\bar{x}_k) \leq 0.$$

Hence, there is at least one weak limiting distribution $\bar{\mu}_\infty$ of the sequence of empirical measures $\bar{\mu}_n := \frac{1}{n} \sum_{0 \leq k \leq n-1} \delta_{\bar{x}_k}$ which is supported by the closed set $\{\varphi = 0\} \subset \{0, 1\} \cup \{f_0 = f_1\}$; on the other $\text{supp}(\mu_\infty)$ is contained in the set $\mathcal{X}_\infty$ of the limiting values of the

sequence $(\bar{x}_n)$ itself (in fact $\bar{\mathcal{X}}_\infty$ is an interval since $(\bar{x}_n)_n$ is bounded and $\bar{x}_{n+1} - \bar{x}_n \rightarrow 0$). Hence $\bar{\mathcal{X}}_\infty \cap (\{0, 1\} \cup \{f_0 = f_1\}) \neq \emptyset$.

If the greedy strategy $(\bar{x}_n)_n$ is optimal then $\text{dist}(\bar{x}_n, \text{argmax } q) \rightarrow 0$ as $n \rightarrow \infty$ *i.e.* $\bar{\mathcal{X}}_\infty \subset \text{argmax } q$. Consequently if

$$\text{argmax } q \cap (\{0, 1\} \cup \{f_0 = f_1\}) = \emptyset \quad (3)$$

then *the purely competitive strategy is never optimal*.

So is the case if

$$f_0(x) = ax \quad \text{and} \quad f_1(x) = b(1 - x), \quad x \in [0, 1],$$

for some positive parameters $a \neq b$, then

$$\text{argmax } q = \{1/2\} \quad \text{and} \quad f_0(1/2) \neq f_1(1/2).$$

In fact, one shows that the greedy strategy $x = (x_n)_{n \geq 1}$ defined by (2) satisfies

$$\bar{x}_n \rightarrow \frac{b}{a+b} \quad \text{and} \quad Q(x) = \frac{ab}{a+b} \quad \text{as} \quad n \rightarrow \infty$$

whereas any optimal (cooperative) strategy (like the *i.i.d.* Bernoulli(1/2) one) yields an asymptotic (relative) global payoff rate

$$Q^* = \max_{[0,1]} q = \frac{a+b}{4}.$$

Note that $Q^* > \frac{ab}{a+b}$ since $a \neq b$. (When $a = b$ the greedy strategy becomes optimal.)

- A more abstract version of Theorem 1 can be established using the same approach. The finite set $\{0, 1, \dots, d\}$ is replaced by a compact metric set K , $\mathcal{P}_{d+1}$ is replaced by the convex set $\mathcal{P}_K$ of probability distributions on K equipped with the weak topology and the continuous function $f : K \times \mathcal{P}_K \rightarrow \mathbb{R}$ still derives from a potential function in some sense.

References

- [1] M. BENAÏM (1999). Dynamics of stochastic algorithms, in J. Azéma et al. eds, *Séminaire de probabilités XXXIII*, L.N. in Math. 1708, 1-68, Springer Verlag, Berlin.
- [2] M. BENAÏM, G. BEN AROUS (2003). A two armed bandit type problem, *Game Theory*, **32**(3), 3-16.